



- *ACHATINELLA APEXFULVA*
- *ACHATINELLA BULIMOIDES*
- ▲ *ACHATINELLA BYRONII/ DECIPIENS*
- ◆ *ACHATINELLA CURTA*
- *ACHATINELLA LILA*
- ▲ *ACHATINELLA LIVIDA*
- ◆ *ACHATINELLA PULCHERRIMA*
- *ACHATINELLA SOWERBYANA*
- *ATYOIDA BISULCATA*
- ▲ *AURICULELLA PULCHRA*
- ◆ *AWAOUS GUAMENSIS*
- *CHASIEMPIS SANDWICHENSIS IBIDIS*
- *LENTIPES CONCOLOR*
- ▲ *MEGALAGRION NIGROHAMATUM NIGROLINEATUM*
- ◆ *PAROREOMYZA MACULATA*
- *VESTIARIA COCCINEA POP 2*



Source: HINHP 2002, CEMML 2002

R:\New\j397-Transformation\GIS\Layouts\Kahuku\_Kawailoa Wildlife Species.mxd - 2/22/04 - YE

Sixteen sensitive wildlife species have been recorded within the Kahuku/Kawailoa Training Areas Region of Influence.

- Legend**
- Drum Road
  - Installation Boundaries
  - - - Region of Influence
  - Roads

## **Sensitive Wildlife Species in the Kahuku /Kawailoa Training Areas Biological Region of Influence**

O'ahu, Hawai'i

**Figure 7-23**